

# Prophesy™

## 0.72G Fungicide on DG Pro



### FEATURES & USE TIPS:

- Systemic fungicide that prevents and controls Dollar Spot, Brown Patch, Summer Patch, rust and other diseases.
- Recommended for control of Necrotic Ring Spot.
- DG PRO™ dispersible carrier - minimizes mower pickup after irrigation.
- Contains propiconazole as the fungicide active ingredient.
- Ideal for use in Snow Mold control programs.
- Convenient, dry-flowable formula.
- Can be applied with rotary or drop spreader.
- Small particle size - ideal for use on golf course greens and low-cut bentgrass fairways.
- Root absorbed for consistent uptake into the turf.

### PRODUCT SPECIFICATIONS:

**Active Ingredient:** 0.72% Propiconazole

*Following is a specimen label that may at some point be revised or otherwise become out of date. It is intended as a guide in providing general information regarding the use of this product. Always read and follow the label on the product container.*

### PRODUCT CODE:

**AGCDGPZ3**

Date: 112807  
Supersedes: 082406  
Label: 17

### PRODUCT CHARACTERISTICS:

**Product Type:**  
Granular Fungicide

**Particle Size:**  
SGN: 100

**Net Weight:**  
40 lbs.

**Coverage:**  
16,000 sq. ft. @ 2.5 lb  
1,000 sq. ft. rate

Prophesy is a trademark of The Andersons, Inc.; DG Pro is a registered trademark of The Andersons, Inc.

*To order, contact your local Andersons Golf Products distributor.*

*For more information, contact Andersons at our website: [www.andersonsgolfproducts.com](http://www.andersonsgolfproducts.com) or call 1-800-225-2639.*



# 0.72G Fungicide on DG<sup>PRO</sup>

Broad spectrum and systemic disease control for turf

**Active Ingredient:**  
 Propiconazole 0.72%

**Other Ingredients:** 99.28%

\*CAS No. 60207-90-1 Total: 100.00%

EPA Reg. No. 9198-219  
 EPA Est. No. 9195-OH-1M, 9198-OH-2B, 9196-AL-001A  
 Undefined letter is first letter used in run code on bag

## KEEP OUT OF REACH OF CHILDREN WARNING/AVISO

Sústed no enterado la etiqueta, busque a alguien para que se la explique a usted en detalle.  
 (If you do not understand the label, find someone to explain it to you in detail.)

### Precautorynary Staments

**Hazards to Humans and Domestic Animals**  
 Can cause eye irritation. Harmful if swallowed, absorbed through skin or inhaled. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wear goggles or face shield. Avoid contact with skin. Avoid breathing dust. Do not get into eyes or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

### Personal Protective Equipment

- Applicators and other handlers must wear:
  - Long-sleeved shirt and long pants
  - Chemical resistant gloves made from any waterproof material
  - Shoes plus socks
  - Goggles or face shield

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### User Safety Recommendations

- Use sprayers before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticides gets inside. Then wash thoroughly and put on clean clothing.

### Environmental Hazards

This pesticide is toxic to fish. Do not apply directly to water, to areas where surface water is present, or to riparian areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters.

### FIRST AID

- If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contacts, if present, after the first 5 minutes; then continue rinsing. Call a poison control center or doctor for treatment advice.
- If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
- If swallowed:** Call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
- If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.

Note to Physician: If inhaled, induce emesis or lavage stomach. Treat symptomatically.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-551-8851 for emergency medical treatment information.

### DIRECTIONS FOR USE

This is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR part 170. This standard contains requirements for the training of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for labeling, documentation, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry areas. The requirements in this box only apply to uses of the product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls, chemical resistant gloves made from any waterproof material, shoes plus socks, protective eyewear.

### NON-AGRICULTURAL USE REQUIREMENTS

The equipment in this box apply to use of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The Worker Protection Standard applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter treated areas until dust has settled.

### Storage and Disposal

Do not store in areas where food or feed is stored or disposed. Bases opened bag by folding top down and securing.  
**PESTICIDE STORAGE:** Store in a clean, dry area or place.  
**PESTICIDE DISPOSAL:** Pesticide wastes may be acutely hazardous. Improper disposal is a violation of Federal law. Pesticide mixtures, or equipment rinse water that cannot be chemically reprocessed must be disposed of according to applicable Federal, state or local procedures. Contact your state pesticide or environmental control agency or the hazardous waste representative at the nearest EPA Regional Office for guidance.  
**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

### General Information

This product is a systemic fungicide for use on turfgrasses for the control:

Dollar spot	Sclerotinia homoeocarpa	Spring dead spot	Leptosphaeria korrae
Brown patch	Rhizoctonia solani		Leptosphaeria tarmani
Red thread	Laetisaria fulviformis		Ophiostoma herpeticum
Pink patch	Limonomyces rosaeppis	Take-all patch	Gaeumannomyces graminis
Powdery mildew	Erysiphe graminis	Leatipot	Bipolaris spp., Drechslera spp.
Strips smut	Ustilago striiformis and Limonomyces rosaeppis	Pink snowmold	Monobachium nivale
Summer patch	Urocystis agropyri	Fusarium patch	Trichia spp.
Yellow patch	Magnaporthe poae	Gray snowmold	Phaeoannellum cereale
	Leptosphaeria korrae	Yellow patch	Phaeoannellum cereale
		Zoysia patch	Phaeoannellum cereale

Do not apply this product through any type of irrigation system.

### Turfgrass and Dichondra Disease Control

- Use this product in a preventative disease control program.
- Properly calibrate spreader to ensure thorough coverage.
- Apply after mowing.
- For control of turf diseases, apply to wettest foliage.
- For control of soil-borne diseases, this product can be watered in after application.
- Optimum turf quality and disease control. Use this product in conjunction with turf management practices that promote good plant health.
- Reduce use of fungicides. Proper diagnosis of the condition causing the disease is important. Use of diagnostic kits or other means of identification of the disease organism is essential to determine the best control measures.
- Do not apply more than 20 lbs. product/1,000 sq. ft. calendar year.

Important: Bermudagrass can be sensitive to this product. Do not exceed 5 lbs./1,000 sq. ft. every 30 days on any variety of bermudagrass. In Florida, do not apply this product to bermudagrass golf greens when temperatures exceed 90°F.

Note: Do not graze animals on treated areas. Do not feed clippings from treated areas to livestock or poultry.

### Turfgrass - Specific Diseases, Rates, and Application Timing

Disease	Lbs. product per 1,000 sq. ft.	Lbs. product per Acre	Application Timing/Interval	Instructions
Dollar spot	2.5	109	14-28 days	Make no more than 3 consecutive applications for dollar spot before rotating to an alternative EPA-registered fungicide having a different mode of action.
Sclerotinia homoeocarpa	2.5	109	14-21 days	Begin applications in May or June before disease is present. Under conditions of high temperatures and high humidity, use this product at shorter intervals.
Brown patch	2.5	109	14-28 days	Apply when conditions are favorable for disease development or when disease is present.
Rhizoctonia solani	2.5	109	14-21 days	Apply when conditions are favorable for disease development.
Powdery mildew	2.5	109	14-28 days	Apply once in the fall after grass becomes dormant and in the early spring before grass starts to grow.
Red thread	2.5	109	14 days	Under light to moderate pressure, apply this product to reduce the severity of leaf spot and melting out caused by Helminthosporium-type pathogens.
Pink patch	2.5	109	14 days	Apply this product beginning in April. Use the 5 lb./1,000 sq. ft. rate on a 2-28 day schedule and the 2.5 lb./1,000 sq. ft. rate on a 14-day schedule.
Summer patch	5	218	28 days	Apply this product to reduce the severity of take-all patch. Make 1-2 fall applications in September and October or when night temperatures drop to 55°F and 1-2 spring applications in April and May, depending on local recommendations.
Take-all patch	2.5-5	109-218	Spring and Fall	Make 1-3 applications. For one application, apply in September or October. For multiple applications, begin applications in August.
Yellow patch	2.5	109	14-28 days	
Zoysia patch	2.5	109	14-28 days	

Disease	Lbs. product per 1,000 sq. ft.	Lbs. product per Acre	Application Timing/Interval	Instructions
Necrotic ring spot	5	218	Fall or spring	Apply in the fall and/or the early spring depending on local recommendations.
Strombold	2.5-5	109-218	Late Fall	Apply one application in the late fall before snow cover. Do not apply on top of snow. For optimum disease control, the 2.5 and 3.75 lb. rates should be used in conjunction with either PCNB or chlorothalonil at label rates.
Gray	2.5-5	109-218	Fall-Early Spring	Apply when conditions are favorable for disease development.
Pink	2.5-5	109-218	Late Fall	Apply one application in the late fall before snow cover. Do not apply on top of snow. Use a 3.75 lb./1,000 sq. ft. rate, use in conjunction with a registered contact fungicide at the label rate.
Fusarium patch	2.5-5	109-218	Early Fall	Make one application in the early fall (mid-September to mid-October) prior to development of disease symptoms. Consult local turfgrass extension experts to determine the optimum application timing in your area.
Microdochium nivale	2.5-5	109-218	14-21 days	Apply when conditions are favorable for disease development.
Yellow patch	2.5	109	14-21 days	
Phaeoannellum cereale	2.5	109	14-21 days	
Zoysia patch	2.5	109	14-21 days	
Phaeoannellum cereale	2.5	109	14-21 days	
Phaeoannellum cereale	2.5	109	14-21 days	

### Suggested Spreader Settings

These suggested spreader settings are not intended to replace calibration. Please calibrate your spreader before applying product.  
 40 lbs. treats 1 acre at the 5.00 lbs./1,000 sq. ft. rate  
 40 lbs. treats 1/2 acre at the 10.00 lbs./1,000 sq. ft. rate  
 40 lbs. treats 1/3 acre at the 15.00 lbs./1,000 sq. ft. rate  
 40 lbs. treats 1/4 acre at the 20.00 lbs./1,000 sq. ft. rate

SPREADER	GROUND OR PTO SPEED	WIDTH OF COVERAGE	RECOMMENDED RATE per 1,000 sq. ft.	
			2.5# (0.78 lbs. a/acre)	5.00# (1.57 lbs. a/acre)
Angelsen AccuPro 2000, SR-2000 (cone 7)	3 mph	9 feet	J	K 1/4
Andersens SS-2 Drop	3 mph	9 feet	4 1/2	5 1/2
Scotts R-BA (cone 7)	3 mph	9 feet	H 1/2	1 3/4
Lawn Model WTR, WFR, HR, 1250	4.5 mph	30 feet	4	4 3/4
Levy PTO at 450 rpm, Pattern III-A	4.5 mph	16 feet	12	14 1/2
Woon 1/3 Series	4.5 mph	16 feet	12	18

### IMPORTANT: READ BEFORE USE

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. Items are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability.

### WARRANTY DISCLAIMER and LIMITATION OF LIABILITY

To the extent consistent with applicable law, Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. To the extent allowed by state law, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. To the extent allowed by state law, Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

### LEGAL RIGHTS:

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS. YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE JURISDICTION TO STATE JURISDICTION. (APR08106)

Manufactured by:  
 The Andersons Professional Division, Inc.  
 P.O. Box 19  
 Maumee, OH 43537  
 www.andersongolfproducts.com

SGN 100

Net Wt. 40 Lbs. (18.14 kg)

17 (AGCBLKPBLL1B)

AGCDGPZ3