

CLICK HERE TO JUMP TO USE DIRECTIONS



# TRIMEC® SOUTHERN

BROADLEAF HERBICIDE FOR SENSITIVE SOUTHERN GRASSES

*Bermudagrass, Bentgrass, Bluegrass,  
Fescue, Zoysiagrass, Centipedegrass  
& St. Augustinegrass*



**ACTIVE INGREDIENTS:**

Dimethylamine salt of (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid	17.37%
Dimethylamine salt of 2,4-dichlorophenoxyacetic acid	18.74%
Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid)	3.85%
<b>INERT INGREDIENTS:</b>	<b>60.04%</b>

TOTAL 100.00%

**THIS PRODUCT CONTAINS:**

1.32 lbs. (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per gallon or 14.35%.  
 1.44 lbs. 2,4-dichlorophenoxyacetic acid equivalent per gallon or 15.57%.  
 0.30 lb. 3,6-dichloro-o-anisic acid equivalent per gallon or 3.20%.  
 Isomer Specific by AOAC Methods.  
 Contains the single isomer form of Mecoprop-p.  
 TRIMEC® is a registered trademark of PBI/Gordon Corporation.

KEEP OUT OF REACH OF CHILDREN

## DANGER

See below for additional Precautionary Statements and First Aid.

KEEP FROM FREEZING



**READ THE ENTIRE LABEL FIRST.  
OBSERVE ALL PRECAUTIONS AND  
FOLLOW DIRECTIONS CAREFULLY.**

### PRECAUTIONARY STATEMENTS

**Hazards to Humans & Domestic Animals**

**DANGER:** Corrosive. Causes damage to eyes. Do not get into eyes, on skin or clothing. Harmful if swallowed or inhaled. Avoid exposure to spray mist. Wear face shield or goggles when mixing, loading or applying this product. When mixing, loading or applying this product, wear long-sleeved shirt, long pants, socks, shoes, chemical-resistant gloves and eye protection. It is recommended that safety glasses include front, brow and temple protection.

After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

**Engineering Control Statements:**

**Containers over 1 gallon and less than 5 gallons:** Persons engaged in open pouring of this product must also wear coveralls or a chemical-resistant apron.

**Containers over 5 gallons or more:** Do not open-pour product from this container. A mechanical system (such as a probe and pump or spigot) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal.

### FIRST AID

<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>Call a poison control center or doctor immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If inhaled:</b>	<ul style="list-style-type: none"> <li>Move person to fresh air.</li> <li>If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth if possible.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If on skin or clothing:</b>	<ul style="list-style-type: none"> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For additional information in case of emergency, call toll free 1-877-800-5556.	

**ENVIRONMENTAL HAZARDS:** This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater. When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources. Do not apply when weather conditions favor drift from target area.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D and MCPP-p have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D and MCPP-p pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried or dust has settled.

For use on residential and other turf sites excluding sod farms.

### STORAGE & DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**STORAGE:** Store in original container in a locked storage area inaccessible to children and domestic animals. Keep from freezing.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your state Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL: Plastic containers:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. **Metal containers:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

## STORAGE & DISPOSAL OBSERVE CAREFULLY

Avoid drift or spray mist onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not pour spray solutions near these plants. Do not use on lippia or dichondra nor on lawns or turf where desirable clovers are present.

Use only lawn type sprayers. Note: Coarse sprays are less likely to drift. Do not spray roots of trees and ornamentals. Do not exceed specified dosages for any area; be particularly careful within the dripline of tree and other ornamental species.

Do not spray while grass is emerging from dormancy. Do not apply to newly seeded lawns until after the second or third mowing. Do not spray when air temperatures exceed 85°F.

Seed can be sown 3 to 4 weeks after application. Do not contaminate domestic or irrigation waters. Be careful not to overdose centipede-grass or St. Augustinegrass. These species are sensitive to 2,4-D, a component of this product.

After using this product, clean sprayer with soap or detergent and water, and rinse thoroughly before applying other pesticides. Failure to observe the above precautions may result in injury.

Do not apply this product through any type of irrigation system.

## HOW MUCH TO USE AND WHERE

The maximum application rate to turf is 0.6 pounds 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

**Bermudagrass & Bentgrass** — Use 0.75 fluid ounces per 1,000 sq. ft., or use 2.0 pints of product per acre. Do not overdose closely-mowed bentgrass. Read the use precautions and observe carefully. Do not use more than the recommended rate. Bermudagrass and bentgrass are moderately sensitive to 2,4-D.

**Bluegrass, Fescues & Zoysiagrass** — Use 0.75 to 1.1 fluid ounces per 1,000 sq. ft., or use 2.0-3.0 pints of product per acre. These species are tolerant to 2,4-D. However, do not overdose.

**St. Augustinegrass & Centipede-grass** — Use 0.37 to 0.55 fluid ounces per 1,000 sq. ft., or use 1-1.5 pints of product per acre. Repeat after 3 weeks if necessary. Do not overdose. Do not apply this product to St. Augustinegrass or centipede-grass during spring green-up which is the transition period between dormancy and active growth. Do not make applications when ambient temperature is below 50°F or above 90°F, or expected to be within 72 hours of application. Do not make applications when grass is under stress from drought, insects, or other herbicides. St. Augustinegrass and centipede-grass are sensitive to 2,4-D and great care should be taken to prevent overdosing. Cultivars vary in tolerance to this product. Do not apply this product to 'Floritam' St. Augustinegrass.

**GENERAL:** Mix in 5 to 300 gallons of water per acre. Use higher rates when weeds are dense or in flower. If hardy weeds persist, make a second application after three weeks. Apply when weeds are young and actively growing. Avoid applications during long excessively dry or hot periods. When grasses are under stress use reduced rates to reduce potential injury. Do not irrigate within 24 hours of application. Do not apply to newly seeded turf until after the second or third mowing. Seed can be sown 3 to 4 weeks after application at recommended rates.

## WEEDS CONTROLLED

Annual fleabane	Cinquefoil	Field oxeye-daisy
Aster, white heath & white prairie	Clover	(*creeping oxeye)
Bedstraw	Cocklebur	Field pennycress
Beggarticks	Compassplant	Filaree, whitestem & redstem
Beggarweed, creeping	Curly dock	Florida pusley
Bindweed	Dandelion	Ground ivy
Birdsfoot trefoil	Dayflower	Groundsel
Black medic	Deadnettle	Hairy bittercress
Broadleaf plantain	Dock	Hawkweed
Buckhorn plantain	Dogfennel	Healall
Bull thistle	Dovefoot geranium	Henbit
Burclover	English daisy	Horsenettle
Burdock, common	False dandelion	Horseweed
Buttercup, creeping	(*spotted catsear & common catsear)	Innocence (Blue-eyed Mary)
Carolina geranium	Field bindweed	Jimsonweed
Carpetweed	(*mornningglory & creeping jenny)	Kochia
Chickweed, common	Field madder	(cont. on next column)
Chicory		

## WEEDS CONTROLLED (cont.)

Lambsquarters	Poison ivy	Virginia-creeper
Lawn burweed	Poison oak	Western salsify
Lespedeza, common	Prostrate knotweed	White clover
Mallow, common	(*knotweed)	(*Dutch clover, honeysuckle clover, white trefoil & purplewort)
Matchweed	Puncturevine	Wild carrot
Mouseear chickweed	Purple cudweed	Wild garlic
Mustard	Purslane	Wild geranium
Nettle	Ragweed	Wild lettuce
Old world diamond flower	Redweed	Wild mustard
<i>Oxalis</i>	Red sorrel	Wild onion
(*yellow woodsorrel & creeping woodsorrel)	(*sheep sorrel)	Wild strawberry
Parsley-piert	Roundleaf greenbriar	Wild violet
Pennsylvania smartweed	Shepherdspurse	Yarrow
Pennywort	Spotted spurge	Yellow rocket
(*dollarweed)	Spurge	and many other broadleaf weeds
Pepperweed	Sunflower	
Pigweed	Thistle	
Pineappleweed	Velvetleaf	
Plantain	(*buttonweed)	
	Venice mallow	
	<i>Veronica</i>	
	(*corn speedwell)	
	Virginia buttonweed	*Synonyms

## LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

**THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED.** This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

896 AP030504  
EPA REG. NO. 2217-655

Manufactured By  
**G pbi/Gordon**  
**CORPORATION**  
An Employee-Owned Company  
1217 West 12th Street  
Kansas City, Missouri 64101  
[www.pbigordon.com](http://www.pbigordon.com)