



BLUE COLORANT

NON-STAINING SPRAY PATTERN INDICATOR AND FERTILIZER INJECTOR TRACER

Inert with most pesticides and fertilizers. Will not permanently stain skin, clothing or equipment. Color is temporary and dissipates with rain and sunlight. Cleans with soap and water.

CAUTION:
KEEP OUT OF REACH OF CHILDREN
DIRECTIONS FOR USE

HAND SPRAYERS:

Use 1/4 to 3/4 oz. (2 to 4 teaspoonfuls) per gallon of water or 1 to 3 oz. per 4 gallons of water.

POWER SPRAY EQUIPMENT:

For large area broadcast applicators use the amount given in the following table:

RATES SA-50 BLUE COLORANT			
GALLONS OF WATER PER ACRE	PER 100 GALLONS OF SPRAY		
	LOW	MEDIUM	HIGH
25	1 QT.	2 QT.	4 QT.
50	1 PT.	1 QT.	2 QT.
100	8 oz.	1 PT.	1 QT.

(Rate approximately equivalent to 1/4 to 3/4 oz. per 1000 sq. ft.)

GREENHOUSE FERTILIZER INJECTORS:

Use the amounts of SA-50 BLUE COLORANT in injector concentrate tank as given below:

INJECTOR RATIO	Amount Per Gallon	Amount Per 10 Gallons	Amount Per 100 Gallons
1:300	1.5 oz.	15 oz.	5 QT.
1:200	1 oz.	10 oz.	3 QT.
1:150	.75 oz.	7.5 oz.	2.33 QT.
1:128	.66 oz.	6.5 oz.	2 QT.
1:100	.5 oz.	5 oz.	3 PT.
1:50	.25 oz. (1 TBS)	2.5 oz.	1.5 PT.

Rates given in both spray application and injector sections are intended as an initial application guide only. Greater or lesser amounts may be indicated depending on color or density of vegetation or by preference of the operator.

DO NOT USE on edible crops. This product is intended for use only on ornamental turf grass (non-grazed), non-crop land, and ornamental plants.

NON-WARRANTY: Seller makes no warranty, expressed or implied as to the use of this material. Buyer assumes all risk and liability for results obtained by use or handling of this material whether or not in accordance with the directions or claimed so to be.

FIRST AID INFORMATION: May be harmful if swallowed. Avoid breathing spray mist. May cause temporary eye and skin irritation. In case of eye contact, flush eyes with plenty of water. In case of skin contact, wash thoroughly with soap and water. If swallowed or if irritation should persist, get medical attention.

STORAGE AND DISPOSAL: Do not reuse container. Rinse thoroughly before disposal. Small spills may be washed away with water. Large spills should be contained and salvaged.

Southern Agricultural Insecticides, Inc.

Palmetto, FL 34220

Hendersonville, NC 28793

Boone, NC 28607

revised 1998